

C(3)).  $^{13}\text{C}$ -NMR: 170.16 (*s*); 137.28 (*d*); 132.24 (*d*); 79.31 (*d*); 78.14 (*d*); 77.94 (*d*); 67.91 (*t*); 60.94 (*t*); 33.15 (*t*); 14.13 (*q*). Anal. calc. for  $\text{C}_{10}\text{H}_{14}\text{O}_4$  (198.22): C 60.59, H 7.12; found: C 60.56, H 7.05.

*Ethyl (RS)-[(1RS,2RS,4RS)-(7-Oxabicyclo[2.2.1]hept-5-en-2-yl)oxy](phenylselanyl)acetate (( $\pm$ )-12).* To a cooled ( $-78^\circ$ ) soln. of ( $\pm$ )-11 (2.0 g, 10.1 mmol) in THF (30 ml) was added under  $\text{N}_2$  a 0.7M LiHMDS soln. in THF (28.9 ml, 20.2 mmol). The mixture was stirred for 5 min and a soln. of diphenyldiselenide (3.15 g, 10.1 mmol) in THF (10 ml) was added. The mixture was allowed to warm to r.t. over 4 h and poured into a mixture of  $\text{Et}_2\text{O}$  (100 ml) and a 1M  $\text{NH}_4\text{Cl}$  soln. (50 ml). The org. layer was washed with brine (50 ml) and dried ( $\text{MgSO}_4$ ). Evaporation of the solvent and FC (AcOEt/PE 1:4) of the crude mixture gave ( $\pm$ )-12 (2.6 g, 73%) as a 1:1 mixture of isomers. EI-MS: 354 (0.99, [M+1] $^+$ ), 243 (10), 197 (7), 157 (16), 129 (9), 111 (9), 95 (38), 91 (11), 77 (52), 67 (100), 55 (27), 51 (26).  $^1\text{H}$ -NMR: 7.68–7.56 (*m*, 2 arom. H); 7.40–7.22 (*m*, 3 arom. H); 6.50 (*dd*, *J* = 6.0, 2.0, H-C(5)); 6.30 (*dd*, *J* = 6.0, 1.8, H-C(6)); 5.41 (*s*, CH-COOEt); 5.02 (*dd*, *J* = 5.0, 2.0 H-C(4)); 4.53 (*ddd*, *J* = 8.0, 4.5, 2.5, H-C(2)); 4.13 (*q*, *J* = 6.5, OCH<sub>2</sub>CH<sub>3</sub>); 2.09 (*ddd*, *J* = 8.0, 5.0, *J* = 12.0, H<sub>exo</sub>-C(3)); 1.21 (*t*, *J* = 6.5, OCH<sub>2</sub>CH<sub>3</sub>); 1.07 (*dd*, *J* = 2.5, *J* = 12.0, H<sub>endo</sub>-C(3)). Anal. calc. for  $\text{C}_{16}\text{H}_{18}\text{O}_4\text{Se}$  (353.27): C 54.40, H 5.14, Se 22.35; found: C 54.30, H 5.12, Se 22.32.

sunlamp for 3 h. Evaporation of the solvent and FC (AcOEt/PE 1:4) of the crude mixture gave ( $\pm$ )-13 (580 mg, 95%) as colorless oil. IR (film): 2980s, 2940s, 1740s, 1580m, 1480m, 1260s, 1200s, 1160m, 1080s, 1040s, 1020s, 980m, 890m, 750s. EI-MS: 354 (26 [M+1] $^+$ ), 353 (2, M $^+$ ), 281 (6), 197 (7), 171 (9), 123 (13), 105 (4), 99 (6), 95 (55), 81 (60), 77 (51), 67 (100), 55 (47).  $^1\text{H}$ -NMR: 7.59–7.49 (*m*, 2 arom. H); 7.32–7.21 (*m*, 3 arom. H); 5.11 (*t*, *J* = 5.0, H-C(7)); 4.58 (*dd*, *J* = 7.0, 5.0, H-C(3)); 4.41 (*d*, *J* = 5.0, H-C(1)); 4.40 (*s*, H-C(5)); 4.13 (*q*, *J* = 7.0, OCH<sub>2</sub>CH<sub>3</sub>); 3.31 (*d*, *J* = 2.0, H-C(5)); 2.63 (*dd*, *J* = 5.0, 2.0, H-C(6)); 1.88 (*ddd*, *J* = 7.5, 5.0, *J* = 12.5, H<sub>exo</sub>-C(2)); 1.53 (*d*, *J* = 12.5, H<sub>endo</sub>-C(2)); 1.22 (*t*, *J* = 7.0, OCH<sub>2</sub>CH<sub>3</sub>).  $^{13}\text{C}$ -NMR: 171.51 (*s*); 133.72 (*d*); 129.14 (*d*); 127.61 (8 *d*); 81.80 (*d*); 79.99 (*d*); 79.38 (*d*); 77.72 (*d*); 61.12 (*t*); 51.27 (*d*); 49.85 (*d*); 40.72 (*t*); 13.93 (*q*); one aromatic C(*d*) is ‘missing’. Anal. calc. for  $\text{C}_{16}\text{H}_{18}\text{O}_4\text{Se}$  (353.27): C 54.40, H 5.14, Se 22.35; found: C 54.30, H 5.12, Se 22.32.

We thank the Swiss National Science Foundation and the Herbette Foundation for financial support. P. R. thanks also the Stiftung für Stipendien auf dem Gebiete der Chemie for the Alfred Werner stipend. We are very grateful to Dr. Yannick Landais for helpful discussions.

Received: July 6, 1994

- [1] Procedures using chiral auxiliaries: a) E. Vieira, P. Vogel, *Helv. Chim. Acta* **1983**, 66, 1865; b) K.A. Black, P. Vogel, *ibid.* **1984**, 67, 1612; c) A. Warm, P. Vogel, *ibid.* **1987**, 70, 690; d) J.-L. Reymond, P. Vogel, *Tetrahedron Lett.*

- hedron: Asymmetry* **1990**, 1, 729. Catalytic enantioselective process: e) E.J. Corey, T.-P. Loh, *Tetrahedron Lett.* **1993**, 34, 3979. [2] P. Vogel, D. Fattori, F. Gasparini, C. Le Drian, *Synlett* **1990**, 173. [3] P. Renaud, J.-P. Vionnet, *J. Org. Chem.* **1993**, 58, 5895. [4] J.-P. Vionnet, K. Schenk, P. Renaud, *Helv. Chim. Acta* **1993**, 76, 2490. [5] P. Renaud, J.-P. Vionnet, *Tetrahedron Lett.* **1991**, 32, 3491. [6] S.M. Waraszkiewicz, H.H. Sun, K.L. Erickson, J. Finer, J. Clardy, *J. Org. Chem.* **1978**, 43, 3194; C.L.D. Jennings-White, A.B. Holmes, P.R. Raithby, *J. Chem. Soc., Chem. Commun.* **1979**, 542; A.B. Holmes, C.L.D. Jennings-White, D.A. Kendrick, *J. Chem. Soc., Chem. Commun.* **1983**, 415. [7] Calculations have shown that a phenylsulfonyl group destabilizes an alkyl radical by 0.8 kcal/mol: R. Merenyi, Z. Janousek, H.G. Viehe, in ‘Substituent Effects in Radical Chemistry’, Eds. H.G. Viehe, Z. Janousek, R. Merenyi, Reidel, Dordrecht, 1986, p. 301. [8] a) J.H. Byers, G.C. Lane, *Tetrahedron Lett.* **1990**, 5697; b) J.H. Byers, T.G. Gleason, K.S. Knight, *J. Chem. Soc., Chem. Commun.* **1991**, 354; c) J.H. Byers, B.C. Harper, *Tetrahedron Lett.* **1992**, 33, 6953; d) J.H. Byers, G.C. Lane, *J. Org. Chem.* **1993**, 58, 3355. [9] D.P. Curran, A.A. Martinesker, S.B. Ko, M. Newcomb, *J. Org. Chem.* **1993**, 58, 4691. [10] G.C. Cox, J.J. Kulagowski, C.J. Moody, E.-R. Sie, *Synlett* **1992**, 975; C.J. Moody, M.J. Davies, R.J. Taylor, *J. Chem. Soc., Perkin Trans. I* **1991**, 1; H.B. Wood, H.-P. Buser, B. Ganem, *J. Org. Chem.* **1992**, 57, 178.

## Korrekturen und Nachträge/ Corrections and Additions

*Chimia* **1994**, 48 (Nr. 9), 444.  
Rolf Steiger, ‘Surface Chemistry of Silver Halide Microcrystals’: Fig. 5 has to be added.




Fig. 5. Imaging SIMS iodide picture of AgBr microcrystals containing 0.13 mol-% iodide. An iodide ring is predominantly formed along edges and corners of the cubic AgBr crystals (1.24- $\mu\text{m}$  edge-length).

C(3)).  $^{13}\text{C}$ -NMR: 170.16 (*s*); 137.28 (*d*); 132.24 (*d*); 79.31 (*d*); 78.14 (*d*); 77.94 (*d*); 67.91 (*t*); 60.94 (*t*); 33.15 (*t*); 14.13 (*q*). Anal. calc. for  $\text{C}_{10}\text{H}_{14}\text{O}_4$  (198.22): C 60.59, H 7.12; found: C 60.56, H 7.05.

*Ethyl (RS)-[(1RS,2RS,4RS)-(7-Oxabicyclo[2.2.1]hept-5-en-2-yl)oxy](phenylselanyl)acetate (( $\pm$ )-12).* To a cooled ( $-78^\circ$ ) soln. of ( $\pm$ )-11 (2.0 g, 10.1 mmol) in THF (30 ml) was added under  $\text{N}_2$  a 0.7M LiHMDS soln. in THF (28.9 ml, 20.2 mmol). The mixture was stirred for 5 min and a soln. of diphenyldiselenide (3.15 g, 10.1 mmol) in THF (10 ml) was added. The mixture was allowed to warm to r.t. over 4 h and poured into a mixture of  $\text{Et}_2\text{O}$  (100 ml) and a 1M  $\text{NH}_4\text{Cl}$  soln. (50 ml). The org. layer was washed with brine (50 ml) and dried ( $\text{MgSO}_4$ ). Evaporation of the solvent and FC (AcOEt/PE 1:4) of the crude mixture gave ( $\pm$ )-12 (2.6 g, 73%) as a 1:1 mixture of isomers. EI-MS: 354 (0.99, [M+1] $^+$ ), 243 (10), 197 (7), 157 (16), 129 (9), 111 (9), 95 (38), 91 (11), 77 (52), 67 (100), 55 (27), 51 (26).  $^1\text{H}$ -NMR: 7.68–7.56 (*m*, 2 arom. H); 7.40–7.22 (*m*, 3 arom. H); 6.50 (*dd*, *J* = 6.0, 2.0, H-C(5)); 6.30 (*dd*, *J* = 6.0, 1.8, H-C(6)); 5.41 (*s*, CH-COOEt); 5.02 (*dd*, *J* = 5.0, 2.0 H-C(4)); 4.53 (*ddd*, *J* = 8.0, 4.5, 2.5, H-C(2)); 4.13 (*q*, *J* = 6.5, OCH<sub>2</sub>CH<sub>3</sub>); 2.09 (*ddd*, *J* = 8.0, 5.0, *J* = 12.0, H<sub>exo</sub>-C(3)); 1.21 (*t*, *J* = 6.5, OCH<sub>2</sub>CH<sub>3</sub>); 1.07 (*dd*, *J* = 2.5, *J* = 12.0, H<sub>endo</sub>-C(3)). Anal. calc. for  $\text{C}_{16}\text{H}_{18}\text{O}_4\text{Se}$  (353.27): C 54.40, H 5.14, Se 22.35; found: C 54.30, H 5.12, Se 22.32.

sunlamp for 3 h. Evaporation of the solvent and FC (AcOEt/PE 1:4) of the crude mixture gave ( $\pm$ )-13 (580 mg, 95%) as colorless oil. IR (film): 2980s, 2940s, 1740s, 1580m, 1480m, 1260s, 1200s, 1160m, 1080s, 1040s, 1020s, 980m, 890m, 750s. EI-MS: 354 (26 [M+1] $^+$ ), 353 (2, M $^+$ ), 281 (6), 197 (7), 171 (9), 123 (13), 105 (4), 99 (6), 95 (55), 81 (60), 77 (51), 67 (100), 55 (47).  $^1\text{H}$ -NMR: 7.59–7.49 (*m*, 2 arom. H); 7.32–7.21 (*m*, 3 arom. H); 5.11 (*t*, *J* = 5.0, H-C(7)); 4.58 (*dd*, *J* = 7.0, 5.0, H-C(3)); 4.41 (*d*, *J* = 5.0, H-C(1)); 4.40 (*s*, H-C(5)); 4.13 (*q*, *J* = 7.0, OCH<sub>2</sub>CH<sub>3</sub>); 3.31 (*d*, *J* = 2.0, H-C(5)); 2.63 (*dd*, *J* = 5.0, 2.0, H-C(6)); 1.88 (*ddd*, *J* = 7.5, 5.0, *J* = 12.5, H<sub>exo</sub>-C(2)); 1.53 (*d*, *J* = 12.5, H<sub>endo</sub>-C(2)); 1.22 (*t*, *J* = 7.0, OCH<sub>2</sub>CH<sub>3</sub>).  $^{13}\text{C}$ -NMR: 171.51 (*s*); 133.72 (*d*); 129.14 (*d*); 127.61 (8 *d*); 81.80 (*d*); 79.99 (*d*); 79.38 (*d*); 77.72 (*d*); 61.12 (*t*); 51.27 (*d*); 49.85 (*d*); 40.72 (*t*); 13.93 (*q*); one aromatic C(*d*) is ‘missing’. Anal. calc. for  $\text{C}_{16}\text{H}_{18}\text{O}_4\text{Se}$  (353.27): C 54.40, H 5.14, Se 22.35; found: C 54.30, H 5.12, Se 22.32.

We thank the Swiss National Science Foundation and the Herbette Foundation for financial support. P. R. thanks also the Stiftung für Stipendien auf dem Gebiete der Chemie for the Alfred Werner stipend. We are very grateful to Dr. Yannick Landais for helpful discussions.

Received: July 6, 1994

- [1] Procedures using chiral auxiliaries: a) E. Vieira, P. Vogel, *Helv. Chim. Acta* **1983**, 66, 1865; b) K.A. Black, P. Vogel, *ibid.* **1984**, 67, 1612; c) A. Warm, P. Vogel, *ibid.* **1987**, 70, 690; d) J.-L. Reymond, P. Vogel, *Tetrahedron Lett.*

- hedron: Asymmetry* **1990**, 1, 729. Catalytic enantioselective process: e) E.J. Corey, T.-P. Loh, *Tetrahedron Lett.* **1993**, 34, 3979. [2] P. Vogel, D. Fattori, F. Gasparini, C. Le Drian, *Synlett* **1990**, 173. [3] P. Renaud, J.-P. Vionnet, *J. Org. Chem.* **1993**, 58, 5895. [4] J.-P. Vionnet, K. Schenk, P. Renaud, *Helv. Chim. Acta* **1993**, 76, 2490. [5] P. Renaud, J.-P. Vionnet, *Tetrahedron Lett.* **1991**, 32, 3491. [6] S.M. Waraszkiewicz, H.H. Sun, K.L. Erickson, J. Finer, J. Clardy, *J. Org. Chem.* **1978**, 43, 3194; C.L.D. Jennings-White, A.B. Holmes, P.R. Raithby, *J. Chem. Soc., Chem. Commun.* **1979**, 542; A.B. Holmes, C.L.D. Jennings-White, D.A. Kendrick, *J. Chem. Soc., Chem. Commun.* **1983**, 415. [7] Calculations have shown that a phenylsulfonyl group destabilizes an alkyl radical by 0.8 kcal/mol: R. Merenyi, Z. Janousek, H.G. Viehe, in ‘Substituent Effects in Radical Chemistry’, Eds. H.G. Viehe, Z. Janousek, R. Merenyi, Reidel, Dordrecht, 1986, p. 301. [8] a) J.H. Byers, G.C. Lane, *Tetrahedron Lett.* **1990**, 5697; b) J.H. Byers, T.G. Gleason, K.S. Knight, *J. Chem. Soc., Chem. Commun.* **1991**, 354; c) J.H. Byers, B.C. Harper, *Tetrahedron Lett.* **1992**, 33, 6953; d) J.H. Byers, G.C. Lane, *J. Org. Chem.* **1993**, 58, 3355. [9] D.P. Curran, A.A. Martinesker, S.B. Ko, M. Newcomb, *J. Org. Chem.* **1993**, 58, 4691. [10] G.C. Cox, J.J. Kulagowski, C.J. Moody, E.-R. Sie, *Synlett* **1992**, 975; C.J. Moody, M.J. Davies, R.J. Taylor, *J. Chem. Soc., Perkin Trans. I* **1991**, 1; H.B. Wood, H.-P. Buser, B. Ganem, *J. Org. Chem.* **1992**, 57, 178.

## Korrekturen und Nachträge/ Corrections and Additions

*Chimia* **1994**, 48 (Nr. 9), 444.  
Rolf Steiger, ‘Surface Chemistry of Silver Halide Microcrystals’: Fig. 5 has to be added.




Fig. 5. Imaging SIMS iodide picture of AgBr microcrystals containing 0.13 mol-% iodide. An iodide ring is predominantly formed along edges and corners of the cubic AgBr crystals (1.24- $\mu\text{m}$  edge-length).